

An Analysis of Bennett's Pebble Game

LANL report LAUR-95-2258

E. Knill, knill@lanl.gov*

May 1995

Abstract

Bennett's pebble game [1, 2] was introduced to obtain better time/space tradeoffs in the simulation of standard Turing machines by reversible ones. So far only upper bounds for the tradeoff based on the pebble game have been published. Here we give a recursion for the time optimal solution of the pebble game given a space bound. We analyze the recursion to obtain an explicit asymptotic expression for the best time-space product.

1 Introduction

In Bennett's pebble game, we have a board of n squares labeled $s_1, \dots, s_n$. Each square may be empty or contain one pebble. A move in the game consists of adding or removing a pebble. A pebble can be added to or removed from square s_i if $i = 1$ or if there is a pebble on square s_{i-1} . Initially, the board is empty. The desired final configuration has exactly one pebble on the board on square s_n . For the motivation and application to reversible computing, we refer the reader to [1, 2].

The pebble game is trivial without resource constraints. At the i 'th step for $1 \leq i \leq n$ we can place a pebble on square s_i . At the $n + i$ 'th step for $1 \leq i \leq n - 1$ we can remove the pebble on square s_{n-i} . This solves the game in $2n - 1$ steps using n pebbles. The time resource T is the number of steps to solve the game, while the space resource S is the number of pebbles used, that

*Address: MS B265, Los Alamos National Laboratory, Los Alamos, NM 87545. This work was performed under the auspices of the U.S. Department of Energy under Contract No. W-7405-ENG-36.

is the maximum number of pebbles on the board at any time. For any solution $\mathcal{S}$ of the pebble game, let $T(\mathcal{S})$ be the number of steps and $S(\mathcal{S})$ the number of pebbles used. Our goal is to determine

$$F(n, S) = \min(T(\mathcal{S}) \mid S(\mathcal{S}) \leq S).$$

Levine and Sherman [2] expressed Bennett's [1] upper bound on $F(n, S)$ by a recursion which they were able to solve. We will obtain an exact recursive expression for $F(n, S)$ which can be evaluated explicitly for any desired n and S . An asymptotic expression for the minimum time space product is obtained. A consequence of this analysis is that an upper bound for the minimum time-space product of quantum factoring (see [3]) is $2^{2\sqrt{n}(1+o(1))}n^3$, where n is the number of bits in the number to be factored. We defer discussion of this observation to another note.

2 Analysis of the pebble game

2.1 Solutions of the pebble game

To obtain a recursive expression for $F(n, S)$ it is useful to model solutions $\mathcal{S}$ by representing the residence of each pebble on a square as an interval. Formally, we view $\mathcal{S}$ as a 0 – 1 matrix, also denoted by $\mathcal{S}$, of dimensions $n \times [-\infty, +\infty]$. $\mathcal{S}_{i,j} = 1$ means that there is a pebble on square s_i after step j . For $\mathcal{S}$ to be a valid solution, we need

- (i) Initial configuration: There exists T_b such that for $t \leq T_b$, $\mathcal{S}_{i,t} = 0$.
- (ii) Final configuration: There exists T_e , such that for $t \geq T_e$, $\mathcal{S}_{i,t} = \delta_{n,i}$, with $\delta_{n,i}$ the usual delta function.
- (iii) One move at a time: If $\mathcal{S}_{i,j} \neq \mathcal{S}_{i,j+1}$ then for all $i' \neq i$, $\mathcal{S}_{i',j} = \mathcal{S}_{i',j+1}$.
- (iv) Add constraint: If $\mathcal{S}_{i,j} = 0$ and $\mathcal{S}_{i,j+1} = 1$, then either $i = 1$ or $\mathcal{S}_{i-1,j} = 1$.
- (v) Remove constraint: If $\mathcal{S}_{i,j} = 0$ and $\mathcal{S}_{i,j-1} = 1$, then either $i = 1$ or $\mathcal{S}_{i-1,j-1} = 1$.

The space requirement of $\mathcal{S}$ is given by the maximum number of 1's in any column of $\mathcal{S}$, while the time requirement is the number of inequalities between adjacent elements in the rows.

Consider square s_i . The i 'th row of $\mathcal{S}$ can be viewed as a collection of intervals of 1's, representing the residence of a pebble on s_i . Thus a (*residence*) *interval* $[k, l]$ of s_i satisfies (1) $k \leq l$, (2) $\mathcal{S}_{i,j} = 1$ for $k \leq j \leq l$, (3) $\mathcal{S}_{i,k-1} = 0$ and (4) $\mathcal{S}_{i,l+1} = 0$.

Let $j, j' \geq 0$. If $I = [k, l]$ is an interval of s_i and $J = [k + j, l - j']$ is an interval of s_{i-1} , then J does not contribute anything to the solution. A better solution in terms of time and space is obtained by removing J (by setting the corresponding entries of $\mathcal{S}$ to 0). This allows us to add another constraint:

(vi) If I is an interval of s_i and J is an interval of s_{i-1} , then $J \not\subseteq I$.

In depicting solutions it is convenient to represent the intervals as horizontal lines at a vertical offset, where no two endpoints have the same horizontal offset. See Figure 2.1 for two examples.


2.2 A recursion for $F(n, S)$

Theorem 2.1 $F(n, S)$ is given by

$$F(n, S) = \begin{cases} 1 & \text{for } n = 1 \text{ and } S \geq 1, \\ \infty & \text{for } n \geq 2 \text{ and } S = 1, \\ \min(F(m, S) + F(m, S - 1) + F(n - m, S - 1) \mid 1 \leq m < n) & \text{for } n \geq 2 \text{ and } S \geq 2. \end{cases}$$

Proof. For $n = 1$, the solution is to place a pebble on s_1 in one step. For $n > 1$, at least 2 pebbles are required.

The recursive expression for $F(n, S)$ is obtained by decomposing the solution matrix. The basic idea is shown in Figure 2.1. We begin by proving that $F(n, S) \geq \min(F(m, S) + F(m, S - 1) + F(n - m, S - 1) \mid 1 \leq m < n)$. Let $n > 1$ and $S > 1$. Let $\mathcal{S}$ be a time optimal solution of the pebble game with $S(\mathcal{S}) \leq S$. Let t_b be the least t such that $\mathcal{S}_{1,t} = 1$. We can assume that for $t \geq t_b$, the t 'th column of $\mathcal{S}$ always has at least one 1. If not we can obtain a more time efficient solution by setting all the entries preceding that column to 0. Let t_e be the largest t such that $\mathcal{S}_{1,t} = 1$. Due to the move constraints, for $t > t_e$, $\mathcal{S}_{i,t} = \delta_{n,i}$. Define $m(\mathcal{S}, t) = \min(i \mid \mathcal{S}_{i,t} = 1)$ and let t_m be the maximum t with $t_b \leq t \leq t_e$ for which $m(\mathcal{S}, t)$ achieves its maximum in this range. Let $m = m(\mathcal{S}, t_m)$. Observe that $m < n$, else we can obtain a better solution by setting $\mathcal{S}_{i,t} = 0$ for $i < m$ and $t > t_m$. We can extract three solutions of smaller problems from $\mathcal{S}$.



Let $\mathcal{S}'_1$ be the submatrix of $\mathcal{S}$ consisting of the first m rows between columns t_b and t_m . Let $\mathcal{S}_1$ be the extension of $\mathcal{S}'_1$ to a solution of the pebble game obtained by letting $\mathcal{S}_{1;i,t} = 0$ for $t < t_b$ and $\mathcal{S}_{1;i,t} = \delta_{m,i}$ for $t > t_m$. Then $\mathcal{S}_1$ is a solution of the pebble game with m squares using at most S pebbles.

Let $\mathcal{S}'_2$ be the submatrix of $\mathcal{S}$ consisting of the first m rows between columns t_m and t_e . Let $\mathcal{S}_2$ be $\mathcal{S}'_2$ with the columns in reverse order and extended to a solution as above for $\mathcal{S}'_1$. Then $\mathcal{S}_2$ is a solution of the pebble game with m squares. We claim that it uses at most $S - 1$ pebbles. To see this we show that for $t \geq t_m$, the maximum $i(t)$ for which $\mathcal{S}_{i,t} = 1$ satisfies $i > m$. Suppose that this is not the case. Let t_1 and t_2 be the right endpoint and the left endpoint of the two intervals of s_1 surrounding t_m . Because of the move constraints, $\mathcal{S}_{i,t} = 0$ for $t_1 < t < t_2$ and $i < m$. This implies that the columns between t_1 and t_2 have only one 1. Since $S \geq 2$, we can set $\mathcal{S}_{1,t}$ to 1 for $t_1 < t < t_2$. This reduces the number of steps by 2, contradicting optimality of $\mathcal{S}$. This argument shows that $i(t_m) > m$. If for $t' > t_m$, $i(t') \leq m$, then we can obtain a more efficient solution by setting $\mathcal{S}_{i,t} = 0$ for $i > m$ and $t \leq t'$.

Let $\mathcal{S}_3$ be the matrix consisting of the rows with index $i > m$. This matrix is a solution of the pebble game with $n - m$ squares. Since this solution achieves the final configuration before t_e , and by the choice of m , $\mathcal{S}_3$ can use at most $S - 1$ pebbles.

Since $T(\mathcal{S}) = T(\mathcal{S}_1) + T(\mathcal{S}_2) + T(\mathcal{S}_3)$, the desired inequality follows.

To show that $F(n, S) \leq \min(F(m, S) + F(m, S - 1) + F(n - m, S - 1) \mid 1 \leq m < n)$, we use the decomposition for the lower bound and show that it can be used constructively for an upper bound. Given $\mathcal{S}_1$, $\mathcal{S}_2$ and $\mathcal{S}_3$ with $S(\mathcal{S}_1) \leq S$, $S(\mathcal{S}_2) \leq S - 1$ and $S(\mathcal{S}_3) \leq S - 1$, we construct a solution for the pebble game with n squares by composing the solutions $\mathcal{S}_i$. Let $t_{b,i}$ and $t_{e,i}$ be defined for $\mathcal{S}_i$ as we defined t_b and t_e for $\mathcal{S}$ above (if $m = 1$ or $n - m = 1$, set the corresponding $t_e = t_b + 1$). By shifting the solutions, we can assume $t_{e,1} + 2 = t_{b,3}$ and $t_{e,2} = t_{e,3} + 2$. Define

$$\mathcal{S}_{i,t} = \begin{cases} \mathcal{S}_{1;i,t} & \text{for } i \leq m \text{ and } t \leq t_{e,1} + 1, \\ \mathcal{S}_{3;i-m,t} & \text{for } m < i \leq n, \\ \mathcal{S}_{2;i,2t_{e,2}-t} & \text{for } i \leq m \text{ and } t_{e,3} + 1 < t, \\ 1 & \text{for } i = m \text{ and } t_{b,3} \leq t \leq t_{e,3} + 1, \\ 0 & \text{otherwise.} \end{cases}$$

Then $\mathcal{S}$ is a solution with $T(\mathcal{S}) = \sum_i T(\mathcal{S}_i)$ satisfying the space bound. ■

The recursion of Theorem 2.1 can be used to compute tables of $F(n, S)$.

The appendix contains some Mathematica expressions which compute $F(n, S)$ dynamically. Tables 1 and 2 contain $F(n, S)$ for small values of n and S . The proof of the theorem can be used to easily obtain the optimal strategy for each n and S .

2.3 Properties of $F(n, S)$

Some immediate observations are as follows:

Observation 2.2 (i) $F(n, S) = 2n - 1$ for $S \geq n$.

(ii) For $S \leq S'$, $F(n, S) \geq F(n, S')$.

(iii) For $n \leq n'$, $F(n, S) \leq F(n', S)$.

We next establish the maximum n for which $F(n, S) \leq \infty$.

Theorem 2.3 $F(n, S) < \infty$ iff $n \leq 2^{S-1}$.

Proof. The statement holds for $S = 1$. We proceed by induction on S and n . If $n \leq 2^{S-1}$ and we let $m = \lfloor n/2 \rfloor$, then $m \leq 2^{S-2}$ and $n - m \leq 2^{S-2}$. It follows that $F(m, S - 1)$, $F(n - m, S - 1)$ and $F(m, S)$ are finite by the induction hypothesis. The recursive expression for $F(n, S)$ now implies that $F(n, S)$ is finite.

If $n > 2^{S-1}$ then either $m > 2^{S-2}$ or $n - m > 2^{S-2}$. Hence every term in the minimum of the recursion for $F(n, S)$ is infinite. ■

To obtain explicit bounds on $F(n, S)$ it is convenient to consider the function

$$\Delta(n, S) = \begin{cases} 0 & \text{if } n \leq 0, \\ F(n + 1, S) - F(n, S) & \text{if } n > 0 \text{ and } F(n + 1, S) \leq \infty, \\ \infty & \text{otherwise.} \end{cases}$$

For $n > 1$ and $S > 1$, let $m(n, S)$ be the least minimizer of the recursive expression for $F(n, S)$. Define $m(0, S) = m(1, S) = 0$. Note that $m(n, S) \leq \lfloor n/2 \rfloor$ by the monotonicity properties of $F(n, S)$.

Lemma 2.4 . For $m = m(n, S)$, $n \geq 1$ and $S \geq 2$,

$$\begin{aligned} \Delta(n - m, S - 1) &> \Delta(m - 1, S) + \Delta(m - 1, S - 1), \\ \Delta(n - m - 1, S - 1) &\leq \Delta(m, S) + \Delta(m, S - 1). \end{aligned}$$

$n \setminus S$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
2	∞	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
3	∞	∞	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
4	∞	∞	9	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
5	∞	∞	∞	11	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9
6	∞	∞	∞	15	13	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11
7	∞	∞	∞	19	17	15	13	13	13	13	13	13	13	13	13	13	13	13	13	13
8	∞	∞	∞	25	21	19	17	15	15	15	15	15	15	15	15	15	15	15	15	15
9	∞	∞	∞	∞	25	23	21	19	17	17	17	17	17	17	17	17	17	17	17	17
10	∞	∞	∞	∞	29	27	25	23	21	19	19	19	19	19	19	19	19	19	19	19
11	∞	∞	∞	∞	33	31	29	27	25	23	21	21	21	21	21	21	21	21	21	21
12	∞	∞	∞	∞	39	35	33	31	29	27	25	23	23	23	23	23	23	23	23	23
13	∞	∞	∞	∞	45	39	37	35	33	31	29	27	25	25	25	25	25	25	25	25
14	∞	∞	∞	∞	53	43	41	39	37	35	33	31	29	27	27	27	27	27	27	27
15	∞	∞	∞	∞	61	47	45	43	41	39	37	35	33	31	29	29	29	29	29	29
16	∞	∞	∞	∞	71	51	49	47	45	43	41	39	37	35	33	31	31	31	31	31
17	∞	∞	∞	∞	∞	57	53	51	49	47	45	43	41	39	37	35	33	33	33	33
18	∞	∞	∞	∞	∞	63	57	55	53	51	49	47	45	43	41	39	37	35	35	35
19	∞	∞	∞	∞	∞	69	61	59	57	55	53	51	49	47	45	43	41	39	37	37
20	∞	∞	∞	∞	∞	77	65	63	61	59	57	55	53	51	49	47	45	43	41	39
21	∞	∞	∞	∞	∞	85	69	67	65	63	61	59	57	55	53	51	49	47	45	43
22	∞	∞	∞	∞	∞	93	73	71	69	67	65	63	61	59	57	55	53	51	49	47
23	∞	∞	∞	∞	∞	101	79	75	73	71	69	67	65	63	61	59	57	55	53	51
24	∞	∞	∞	∞	∞	109	85	79	77	75	73	71	69	67	65	63	61	59	57	55
25	∞	∞	∞	∞	∞	117	91	83	81	79	77	75	73	71	69	67	65	63	61	59
26	∞	∞	∞	∞	∞	125	97	87	85	83	81	79	77	75	73	71	69	67	65	63
27	∞	∞	∞	∞	∞	135	105	91	89	87	85	83	81	79	77	75	73	71	69	67
28	∞	∞	∞	∞	∞	145	113	95	93	91	89	87	85	83	81	79	77	75	73	71
29	∞	∞	∞	∞	∞	155	121	99	97	95	93	91	89	87	85	83	81	79	77	75
30	∞	∞	∞	∞	∞	167	129	105	101	99	97	95	93	91	89	87	85	83	81	79
31	∞	∞	∞	∞	∞	179	137	111	105	103	101	99	97	95	93	91	89	87	85	83
32	∞	∞	∞	∞	∞	193	145	117	109	107	105	103	101	99	97	95	93	91	89	87
33	∞	∞	∞	∞	∞	∞	153	123	113	111	109	107	105	103	101	99	97	95	93	91
34	∞	∞	∞	∞	∞	∞	161	129	117	115	113	111	109	107	105	103	101	99	97	95
35	∞	∞	∞	∞	∞	∞	169	137	121	119	117	115	113	111	109	107	105	103	101	99
36	∞	∞	∞	∞	∞	∞	177	145	125	123	121	119	117	115	113	111	109	107	105	103
37	∞	∞	∞	∞	∞	∞	185	153	129	127	125	123	121	119	117	115	113	111	109	107
38	∞	∞	∞	∞	∞	∞	193	161	135	131	129	127	125	123	121	119	117	115	113	111
39	∞	∞	∞	∞	∞	∞	201	169	141	135	133	131	129	127	125	123	121	119	117	115
40	∞	∞	∞	∞	∞	∞	209	177	147	139	137	135	133	131	129	127	125	123	121	119
41	∞	∞	∞	∞	∞	∞	217	185	153	143	141	139	137	135	133	131	129	127	125	123
42	∞	∞	∞	∞	∞	∞	225	193	159	147	145	143	141	139	137	135	133	131	129	127
43	∞	∞	∞	∞	∞	∞	235	201	165	151	149	147	145	143	141	139	137	135	133	131
44	∞	∞	∞	∞	∞	∞	245	209	173	155	153	151	149	147	145	143	141	139	137	135
45	∞	∞	∞	∞	∞	∞	255	217	181	159	157	155	153	151	149	147	145	143	141	139
46	∞	∞	∞	∞	∞	∞	265	225	189	163	161	159	157	155	153	151	149	147	145	143
47	∞	∞	∞	∞	∞	∞	275	233	197	169	165	163	161	159	157	155	153	151	149	147
48	∞	∞	∞	∞	∞	∞	285	241	205	175	169	167	165	163	161	159	157	155	153	151
49	∞	∞	∞	∞	∞	∞	297	249	213	181	173	171	169	167	165	163	161	159	157	155
50	∞	∞	∞	∞	∞	∞	309	257	221	187	177	175	173	171	169	167	165	163	161	159

Table 1. Table of $F(n, S)$ for $1 \leq n \leq 50$ and $1 \leq S \leq 20$.

$n \setminus S$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
51	∞	∞	∞	∞	∞	∞	321	265	229	193	181	179	177	175	173	171	169	167	165	163
52	∞	∞	∞	∞	∞	∞	333	273	237	199	185	183	181	179	177	175	173	171	169	167
53	∞	∞	∞	∞	∞	∞	345	281	245	205	189	187	185	183	181	179	177	175	173	171
54	∞	∞	∞	∞	∞	∞	359	289	253	213	193	191	189	187	185	183	181	179	177	175
55	∞	∞	∞	∞	∞	∞	373	297	261	221	197	195	193	191	189	187	185	183	181	179
56	∞	∞	∞	∞	∞	∞	387	305	269	229	201	199	197	195	193	191	189	187	185	183
57	∞	∞	∞	∞	∞	∞	401	313	277	237	207	203	201	199	197	195	193	191	189	187
58	∞	∞	∞	∞	∞	∞	415	321	285	245	213	207	205	203	201	199	197	195	193	191
59	∞	∞	∞	∞	∞	∞	433	329	293	253	219	211	209	207	205	203	201	199	197	195
60	∞	∞	∞	∞	∞	∞	451	337	301	261	225	215	213	211	209	207	205	203	201	199
61	∞	∞	∞	∞	∞	∞	469	345	309	269	231	219	217	215	213	211	209	207	205	203
62	∞	∞	∞	∞	∞	∞	489	353	317	277	237	223	221	219	217	215	213	211	209	207
63	∞	∞	∞	∞	∞	∞	509	361	325	285	243	227	225	223	221	219	217	215	213	211
64	∞	∞	∞	∞	∞	∞	531	369	333	293	249	231	229	227	225	223	221	219	217	215
65	∞	∞	∞	∞	∞	∞	∞	379	341	301	257	235	233	231	229	227	225	223	221	219
66	∞	∞	∞	∞	∞	∞	∞	389	349	309	265	239	237	235	233	231	229	227	225	223
67	∞	∞	∞	∞	∞	∞	∞	399	357	317	273	243	241	239	237	235	233	231	229	227
68	∞	∞	∞	∞	∞	∞	∞	409	365	325	281	249	245	243	241	239	237	235	233	231
69	∞	∞	∞	∞	∞	∞	∞	419	373	333	289	255	249	247	245	243	241	239	237	235
70	∞	∞	∞	∞	∞	∞	∞	429	381	341	297	261	253	251	249	247	245	243	241	239
71	∞	∞	∞	∞	∞	∞	∞	439	389	349	305	267	257	255	253	251	249	247	245	243
72	∞	∞	∞	∞	∞	∞	∞	449	397	357	313	273	261	259	257	255	253	251	249	247
73	∞	∞	∞	∞	∞	∞	∞	459	405	365	321	279	265	263	261	259	257	255	253	251
74	∞	∞	∞	∞	∞	∞	∞	469	413	373	329	285	269	267	265	263	261	259	257	255
75	∞	∞	∞	∞	∞	∞	∞	481	421	381	337	291	273	271	269	267	265	263	261	259
76	∞	∞	∞	∞	∞	∞	∞	493	429	389	345	297	277	275	273	271	269	267	265	263
77	∞	∞	∞	∞	∞	∞	∞	505	437	397	353	305	281	279	277	275	273	271	269	267
78	∞	∞	∞	∞	∞	∞	∞	517	445	405	361	313	285	283	281	279	277	275	273	271
79	∞	∞	∞	∞	∞	∞	∞	529	453	413	369	321	289	287	285	283	281	279	277	275
80	∞	∞	∞	∞	∞	∞	∞	541	461	421	377	329	295	291	289	287	285	283	281	279
81	∞	∞	∞	∞	∞	∞	∞	553	469	429	385	337	301	295	293	291	289	287	285	283
82	∞	∞	∞	∞	∞	∞	∞	565	477	437	393	345	307	299	297	295	293	291	289	287
83	∞	∞	∞	∞	∞	∞	∞	579	485	445	401	353	313	303	301	299	297	295	293	291
84	∞	∞	∞	∞	∞	∞	∞	593	493	453	409	361	319	307	305	303	301	299	297	295
85	∞	∞	∞	∞	∞	∞	∞	607	501	461	417	369	325	311	309	307	305	303	301	299
86	∞	∞	∞	∞	∞	∞	∞	621	509	469	425	377	331	315	313	311	309	307	305	303
87	∞	∞	∞	∞	∞	∞	∞	635	517	477	433	385	337	319	317	315	313	311	309	307
88	∞	∞	∞	∞	∞	∞	∞	649	525	485	441	393	343	323	321	319	317	315	313	311
89	∞	∞	∞	∞	∞	∞	∞	663	533	493	449	401	349	327	325	323	321	319	317	315
90	∞	∞	∞	∞	∞	∞	∞	677	541	501	457	409	357	331	329	327	325	323	321	319
91	∞	∞	∞	∞	∞	∞	∞	691	549	509	465	417	365	335	333	331	329	327	325	323
92	∞	∞	∞	∞	∞	∞	∞	705	557	517	473	425	373	339	337	335	333	331	329	327
93	∞	∞	∞	∞	∞	∞	∞	721	565	525	481	433	381	345	341	339	337	335	333	331
94	∞	∞	∞	∞	∞	∞	∞	737	575	533	489	441	389	351	345	343	341	339	337	335
95	∞	∞	∞	∞	∞	∞	∞	753	585	541	497	449	397	357	349	347	345	343	341	339
96	∞	∞	∞	∞	∞	∞	∞	769	595	549	505	457	405	363	353	351	349	347	345	343
97	∞	∞	∞	∞	∞	∞	∞	785	605	557	513	465	413	369	357	355	353	351	349	347
98	∞	∞	∞	∞	∞	∞	∞	801	615	565	521	473	421	375	361	359	357	355	353	351
99	∞	∞	∞	∞	∞	∞	∞	817	625	573	529	481	429	381	365	363	361	359	357	355
100	∞	∞	∞	∞	∞	∞	∞	833	635	581	537	489	437	387	369	367	365	363	361	359

Table 2. Table of $F(n, S)$ for $51 \leq n \leq 100$ and $1 \leq S \leq 20$.

Proof. Except for boundary cases, the result is obtained by comparing the expression in the minimum of the recursion of $F(n, S)$ for $m = m(n, S)$, $m = m(n, S) - 1$ and $m = m(n, S) + 1$. The boundary cases are readily checked. ■

Lemma 2.5 *The following hold:*

- (i) $\Delta(n, S)$ is nondecreasing in n .
- (ii) Let $n \geq 1$ and $S \geq 2$. If $\Delta(n - m(n, S), S - 1) \leq \Delta(m(n, S), S) + \Delta(m(n, S), S - 1)$, then $m(n + 1, S) = m(n, S)$. If not, then $m(n + 1, S) = m(n, S) + 1$.
- (iii) Let $n \geq 1$ and $S \geq 2$. Then either $\Delta(n, S) = \Delta(n - m(n, S), S - 1)$ or $\Delta(n, S) = \Delta(m(n, S), S) + \Delta(m(n, S), S - 1)$, whichever is smaller.

Proof. We prove the lemma by induction on S and n . For $S = 1$, $\Delta(n, S)$ is nondecreasing in n by inspection. Let $S > 1$. We have $m(0, S) = m(1, S) = 0$ by definition. Note that $m(2, S) = 1$. $\Delta(0, S) = 0$ and $\Delta(1, S) = 2$ which gives the statement for $n = 0$ and $n = 1$. Let $n \geq 2$. By induction, $\Delta(n', S')$ is nondecreasing for $S' = S - 1$ and for $S' = S$ and $n' < n$. Let $m = m(n, S)$. Using Lemma 2.4 we get for $m' < m$

$$\begin{aligned} \Delta(n + 1 - m' - 1, S - 1) &\geq \Delta(n - m, S - 1) \\ &> \Delta(m - 1, S) + \Delta(m - 1, S - 1) \\ &\geq \Delta(m', S) + \Delta(m', S - 1), \end{aligned}$$

which means that $m' \neq m(n + 1, S)$. Similarly, for $m' > m + 1$,

$$\begin{aligned} \Delta(n + 1 - m', S - 1) &\leq \Delta(n - m - 1, S - 1) \\ &\leq \Delta(m, S) + \Delta(m, S - 1) \\ &\leq \Delta(m' - 1, S) + \Delta(m' - 1, S - 1), \end{aligned}$$

so that $m' \neq m(n + 1, S)$. This leaves only $m(n + 1, S) = m$ or $m(n + 1, S) = m + 1$. The former holds iff $\Delta(n - m, S - 1) \leq \Delta(m, S) + \Delta(m, S - 1)$. In this case, $\Delta(n, S) = \Delta(n - m, S - 1)$ and otherwise $\Delta(n, S) = \Delta(m, S) + \Delta(m, S - 1)$. In the former case, by induction $\Delta(n - 1, S) = \Delta(n - 1 - m, S - 1)$ or $\Delta(n - 1, S) = \Delta(m - 1, S) + \Delta(m - 1, S - 1)$, whichever is smaller, so that by monotonicity, $\Delta(n, S) \geq \Delta(n - 1, S)$. In the latter case, $\Delta(n - 1, S) = \Delta(n - 1 - (m - 1), S - 1)$

or $\Delta(n-1, S) = \Delta(m-2, S) + \Delta(m-2, S-1)$, whichever is smaller, and again it is clear that $\Delta(n, S) \geq \Delta(n-1, S)$. ■

Lemma 2.6 *Let $m = m(n, S)$. Then*

$$\begin{aligned} \Delta(n-m-1, S-1) &\leq \Delta(n, S) \leq \Delta(n-m, S-1), \\ \Delta(m-1, S) + \Delta(m-1, S-1) &\leq \Delta(n, S) \leq \Delta(m, S) + \Delta(m, S-1). \end{aligned}$$

Proof. This is an immediate consequence of Lemmas 2.4 and 2.5. ■

Lemma 2.7 *$\Delta(n, S)$ is nonincreasing in S .*

Proof. The lemma follows from (i) of Lemma 2.5, Lemma 2.6 and by induction. Let $m = m(n, S+1)$ and $m' = m(n, S)$. We have

$$\begin{aligned} \Delta(n, S+1) &= \min(\Delta(n-m, S+1), \Delta(m, S+1) + \Delta(m, S)) \\ &\leq \min(\Delta(n-m, S), \Delta(m, S) + \Delta(m, S-1)). \end{aligned}$$

If $m' = m$, then the last expression is $\Delta(n, S)$ and we are done. If $m' > m$, then by Lemma 2.6,

$$\begin{aligned} \Delta(n, S) &\geq \Delta(m'-1, S) + \Delta(m'-1, S-1) \\ &\geq \Delta(m, S) + \Delta(m, S-1). \end{aligned}$$

Similarly, for $m' < m$,

$$\begin{aligned} \Delta(n, S) &\geq \Delta(n-m'-1, S-1) \\ &\geq \Delta(n-m, S). \end{aligned}$$

In either case, the desired inequality follows. ■

For $k \geq 0$, let $x_{k,S} = \min(n \mid \Delta(n, S) > 2^k)$. This is well defined and nondecreasing in k . We have $x_{0,S} = 1$, $x_{1,S} = S$, and for $k \geq 0$, $x_{k,1} = 1$.

Lemma 2.8 *For $S \geq 2$ and $k \geq 0$ we have*

$$\begin{aligned} x_{k+1,S} &\geq x_{k+1,S-1} + x_{k,S-1}, \\ x_{k+1,S} &\leq x_{k+1,S-1} + x_{k,S}, \\ x_{k+1,S} &\leq 2x_{k+1,S-1}. \end{aligned}$$

Proof. The proof is by induction on S and k . By Lemma 2.6, if either $\Delta(n - m, S - 1) \leq 2^{k+1}$ or $\Delta(m, S - 1) \leq 2^k$, then $\Delta(n, S) \leq 2^{k+1}$. This must occur if $n < x_{k+1, S-1} + x_{k, S-1}$ so that $x_{k+1, S} \geq x_{k+1, S-1} + x_{k, S-1}$.

Suppose that $x_{k, S} > x_{k+1, S-1}$ and $n \geq 2x_{k+1, S-1}$. Since $m = m(n, S) \leq \lfloor n/2 \rfloor$, $n - m \geq x_{k+1, S-1}$. If $m = n - m = x_{k+1, S-1}$, then $\Delta(n - m, S - 1) > 2^{k+1}$ and $\Delta(m, S) + \Delta(m, S - 1) \geq \Delta(m, S - 1) > 2^{k+1}$ which gives $\Delta(n, S) > 2^{k+1}$. If $m < x_{k+1, S-1}$, then $n - m > x_{k+1, S-1}$ and by Lemma 2.6, $2^{k+1} < \Delta(n - m - 1, S - 1) \leq \Delta(n, S)$, so the last inequality follows.

Suppose that $x_{k, S} \leq x_{k+1, S-1}$ and $n \geq x_{k, S} + x_{k+1, S-1}$. If $m = m(n, S) > x_{k, S}$, then the second inequality of Lemma 2.6 and monotonicity of Δ imply $\Delta(n, S) > 2^{k+1}$. If $m < x_{k, S}$, then $n - m > x_{k+1, S-1}$ and the first inequality of Lemma 2.6 can be applied to show that $\Delta(n, S) > 2^{k+1}$. Suppose that $m = x_{k, S}$ and $n = x_{k+1, S-1}$. Then both $\Delta(n - m, S - 1) > 2^{k+1}$ and $\Delta(m, S - 1) \geq \Delta(m, S) > 2^k$ and again $\Delta(n, S) > 2^{k+1}$. This completes the proof of the lemma. $\blacksquare$

Lemma 2.9 For $k \geq 0$ and $S \geq 2$, let

$$\begin{aligned} x_{k, S}^{(l)} &= \sum_{i=0}^k \binom{S-1}{i}, \\ x_{k, S}^{(u)} &= \min \left(\binom{S+k-1}{k}, 2^{S-1} \right). \end{aligned}$$

Then $x_{k, S}^{(l)} \leq x_{k, S} \leq x_{k, S}^{(u)}$.

Proof. It is straightforward to check that $x^{(l)}$ satisfies the recursion $x_{k+1, S}^{(l)} = x_{k+1, S-1}^{(l)} + x_{k, S-1}^{(l)}$ with the initial condition $x_{1, S}^{(l)} = 1$. The function $y_{k, S} = \binom{S+k-1}{k}$ satisfies the recursion $y_{k+1, S} = y_{k+1, S-1} + y_{k, S}$ with the correct initial condition. The bound 2^{S-1} in the definition of $x^{(u)}$ comes from Theorem 2.3. The result follows by Lemma 2.8. $\blacksquare$

Lemma 2.9 cannot be used to obtain “nice” expressions for $F(n, S)$ but yields lower and upper bounds by summing $\Delta(n, S)$.

Lemma 2.10 For $1 \leq k \leq S - 1$ and $S \geq 2$

$$F(x_{k, S}^{(l)}, S) \leq \sum_{i=0}^k \binom{S-1}{i} 2^{i+1},$$

$$F(x_{k,S}^{(u)}, S) \geq \sum_{i=0}^k \binom{S+i-2}{i} (2^i + 1).$$

Proof. The inequalities are obtained by summing $\Delta(n, S)$ and making use of Lemma 2.9, ignoring the second case for $x_{k,S}^{(u)}$. Observe that $\binom{S+k-1}{k} = \sum_{i=0}^k \binom{S+i-2}{i}$. ■

2.4 Asymptotic bounds

For asymptotics where $S = O(\log(n))$, let

$$f(\gamma, S) = \frac{1}{S} \log F(\lfloor 2^{\gamma S} \rfloor, S).$$

All logarithms are base 2. The floor/ceiling notation will be omitted (or absorbed into the ‘ o ’ and ‘ O ’ notation) for real numbers unless needed for clarity. The definition of $f(\gamma, S)$ implies that $F(n, \log(n)/\gamma) = n^{f(\gamma, \log(n)/\gamma)/\gamma + o(1)}$.

For $0 \leq \gamma \leq 1$, let $H(\gamma)$ denote the usual entropy function

$$H(\gamma) = -\gamma \log(\gamma) - (1 - \gamma) \log(1 - \gamma).$$

Theorem 2.11 *Let $\gamma \leq 1/2$. Then*

$$\begin{aligned} f(H(\gamma), S) &\leq \gamma + H(\gamma) + o(1), \\ f((1 + \gamma)H(\gamma/(1 + \gamma)), S) &\geq \gamma + (1 + \gamma)H(\gamma/(1 + \gamma)) + o(1), \end{aligned}$$

where the $o(1)$ are with respect to $S \rightarrow \infty$.

The proof of the theorem is below. First we make a few observations. For small γ , the two expressions in the bounds are the same to within $O(\gamma^2)$.

$$\begin{aligned} H(\gamma) &= -\gamma \log(\gamma) + \log(e)\gamma + O(\gamma^2) \\ &= \gamma(\log(e/\gamma)) + O(\gamma^2), \\ (1 + \gamma)H(\gamma/(1 + \gamma)) &= -\gamma \log(\gamma) + (1 + \gamma) \log(1 + \gamma) \\ &= \gamma(\log(e/\gamma)) + O(\gamma^2). \end{aligned}$$

So for small γ , the bounds are asymptotically tight. The upper bound is an asymptotic version of that given in [2] with a slightly better constant. More

specifically, let $\epsilon = \frac{1}{\log(e/\gamma)}$. Then $\gamma = e2^{-1/\epsilon}$ and $H(\gamma) = (1+\gamma)H(\gamma/(1+\gamma)) + O(\gamma^2) = \frac{\gamma}{\epsilon} + O(\gamma^2)$. Substituting this in the theorem gives

$$F\left(n, \frac{\epsilon}{e} 2^{1/\epsilon} \log(n)\right) = n^{1+\epsilon+2^{-O(\epsilon(1/\epsilon))+o_n(1)}},$$

where we used suffixes on the ‘ O ’ and ‘ o ’ notation to clarify the limit involved.

Proof of Theorem 2.11. A consequence of the Stirling approximation of $n!$ is that for α constant, $\alpha \leq 1/2$,

$$\sum_{i=0}^{\alpha S} \binom{S}{i} = O\left(\binom{S}{\alpha S}\right) = 2^{H(\alpha)S(1+o(1))}.$$

Thus, for $n = 2^{H(\gamma)S}$, $\Delta(n, S) \leq 2^{\gamma S(1+o(1))}$. Using these observations and Lemma 2.10, one can estimate

$$\begin{aligned} F(n, S) &\leq \binom{S}{\gamma S} 2^{\gamma S(1+o(1))} \\ &\leq 2^{(\gamma+H(\gamma))S(1+o(1))}. \end{aligned}$$

Similarly, for α constant,

$$\sum_{i=0}^{\alpha S} \binom{S+i}{i} = O\left(\binom{S(1+\alpha)}{\alpha S}\right) = 2^{(1+\alpha)H(\alpha/(1+\alpha))S(1+o(1))},$$

so that for $n = 2^{(1+\gamma)H(\gamma/(1+\gamma))}$, $\Delta(n, S) \geq 2^{\gamma S(1+o(1))}$. This gives the second inequality. ■

2.5 Asymptotics of TS .

The minimum time-space product is asymptotically determined by the next theorem.

Theorem 2.12

$$TS(n) = 2^{2\sqrt{\log(n)}(1+o(1))} n.$$

Proof. To show that $TS(n) \leq 2^{2\sqrt{\log(n)}(1+o(1))}$, let $S = 2^{\sqrt{\log(n)}}$. Then

$$\begin{aligned} x_{c\sqrt{\log(n)}, S} &\geq \sum_{i=0}^{c\sqrt{\log(n)}} \binom{S}{i} \\ &\geq \binom{2^{\sqrt{\log(n)}}}{c\sqrt{\log(n)}} \\ &\geq n^{c+o(1)}. \end{aligned}$$

This implies that $\Delta(n, S) \leq 2^{\sqrt{\log(n)}(1+o(1))}$ and

$$F(n, S) \leq 2^{\sqrt{\log(n)}(1+o(1))} n.$$

For the lower bound, we estimate $F(n, S)$ for $S = 2^{d\sqrt{\log(n)}}$ from below.

$$\begin{aligned} x_{c\sqrt{\log(n)}, S} &\leq \binom{S + c\sqrt{\log(n)} - 1}{c\sqrt{\log(n)}} \\ &\leq \left(S + c\sqrt{\log(n)} \right)^{c\sqrt{\log(n)}} \\ &\leq 2^{cd\log(n)(1+o(1))} = n^{cd(1+o(1))}. \end{aligned}$$

This implies that $\Delta(n, S) \geq 2^{\frac{\sqrt{\log(n)}}{d}(1+o(1))}$ and

$$\begin{aligned} F(n, S)S &\geq 2^{\sqrt{\log(n)}(d+\frac{1}{d})(1+o(1))} n(1-o(1)) \\ &\geq 2^{2\sqrt{\log(n)}(1+o(1))} n, \end{aligned}$$

since $d + \frac{1}{d} \geq 2$ for all $d > 0$. We used the observation that the last term of the sum $\sum_{i=0}^k \binom{S+i-2}{i}$ asymptotically dominates it. ■

A Appendix

A.1 Mathematica expressions for dynamic evaluation of $F(n, S)$

```
FBen[1,S_] := If[S >= 1, 1, Infinity];
FBen[n_,0] := Infinity;
```

```
FBen[n_,S_] := FBen[n,S] = Module[{m},
  Min[Table[FBen[m,S]+FBen[m,S-1]+FBen[n-m,S-1],{m,1,n-1}]]
];
```

If the goal is to compute a table, then Lemma 2.5 can be used for efficient computation by keeping track of $m(n, S)$.

```
FBenTable[nmax_,Smax_] := Module[{m,n,nt,S,T,Tt,t1,t2,t},
  T = {Join[{1}, Table[Infinity,{n,2,nmax}]]};
  For[S=2, S<=Smax, S++,
    m=1;
```

```

Tt = {1,3};
For[n=3, n<=nmax, n++,
  t1=Tt[[m]]+T[[S-1,m]]+T[[S-1,n-m]];
  t2=Tt[[m+1]]+T[[S-1,m+1]]+T[[S-1,n-m-1]];
  If[t2<t1,
    m++;
    t = t2;
    ,
    t = t1;
  ];
  Tt = Append[Tt,t];
  If[t==Infinity,
    Tt = Join[Tt,Table[Infinity,{nt,n+1,nmax}]];
    Break[];
  ,Null]
];
T = Append[T,Tt];
];
Return[T];
];

```

References

- [1] C. H. Bennett. Time/space trade-offs for reversible computation. *SIAM J. Comput.*, 18:766–776, 1989.
- [2] R. Y. Levine and A. T. Sherman. A note on Bennett’s time-space trade-off for reversible computation. *SIAM J. Comput.*, 19:673–677, 1990.
- [3] P. Shor. Algorithms for quantum computation: Discrete logarithms and factoring. In *Proceedings of the 35’t Annual Symposium on Foundations of Computer Science*, pages 124–134. IEEE Press, 1994.